
Complex charged particle behavior in dusty plasmas and dust crystals: overview of a new test-bed for nonlinear theories

Ioannis KOURAKIS*

Universiteit Gent, Sterrenkundig Observatorium Krijgslaan 281, B-9000 Gent,
Belgium

and

Ruhr Universität Bochum, Fakultät für Physik und Astronomie
Institut für Theoretische Physik IV, D-44780 Bochum, Germany

* Electronic Address: Ioannis.Kourakis@UGent.be, ioannis@tp4.rub.de

Dusty Plasmas (DP) (or *Complex Plasmas*) are large ensembles of interacting particles, consisting of electrons e^- , ions i^+ and massive, heavily charged, mesoscopic-sized defects, i.e. *dust* particulates d^- (or, less often, d^+). The presence of the extra *dust* species has recently been shown to modify the plasma properties substantially and allows for new charged matter configurations, including liquid-like and solid (quasi-*crystalline*) strongly configured phases. These aspects are briefly introduced, at a fundamental level, with emphasis on dust crystal configurations.

The nonlinear aspects of horizontal (longitudinal, acoustic mode) as well as vertical (transverse, optical mode) motion of charged dust grains in a (one-dimensional) dusty plasma monolayer, suspended in a plasma discharge under the combined action of gravity and electrical forces, are discussed. Different types of localized excitations, predicted by nonlinear wave theories, are reviewed and conditions for their occurrence (and characteristics) in DP crystals are discussed. Dust crystals are shown to support nonlinear *kink*-shaped supersonic solitary excitations [1], related to longitudinal (in-plane) dust grain displacement, as well as modulated *envelope localized modes* associated with either longitudinal (in-plane, acoustic) [2] or transverse (off-plane, inverse-optic) [3] oscillations.

Dust crystals consist a highly discrete system, *par excellence*. The occurrence of highly localized *Discrete Breather*-type excitations (Intrinsic Localized Modes) is briefly addressed, from first principles [4, 5, 6, 7]. This possibility opens new directions in nonlinear lattice dynamics- and dusty plasma physics- related research.

The relation to previous results on atomic chains as well as experimental results on strongly-coupled dust layers in gas discharge plasmas is discussed. The relevance of this research to strongly-coupled charged particle configurations, namely Ultra-cold Plasmas (UCPs) confined in Penning trap experiments [8], is pointed out.

-
- [1] I. Kourakis and P. K. Shukla, particle motion in *European Physical Journal D* **29**, 247 (2004).
 - [2] I. Kourakis and P. K. Shukla, *Physics of Plasmas* **11** (4), 1384 (2004).
 - [3] I. Kourakis and P. K. Shukla, *Physics of Plasmas* **11** (5), 2322 (2004).
 - [4] I. Kourakis and P. K. Shukla, *Physics of Plasmas* **12** (1), 014502 (2005).
 - [5] I. Kourakis, V. Basios and P. K. Shukla, *Proc. Int. Conf. Plasma Phys.* (2004), P1-104; online at <http://hal.ccsd.cnrs.fr/ccsd-00001892/en/>.
 - [6] I. Kourakis and P. K. Shukla, *International Journal of Bifurcation and Chaos* **16** (6), to appear (2006).

Abstract for Dynamics Days Crete 2006

Topic: Pattern Formation

- [7] V. Koukouloyannis and I. Kourakis, *Existence of localized modes in systems obeying an inverse dispersion law: the case of transverse dust-lattice vibrations*, in preparation.
- [8] T.C. Killian, *Plasmas put in order*, *Nature* **429**, 815 (2004).